

Energy - Momentum Tensor (p. 49)

(or Stress - Energy tensor) $g = \text{diag}(-1, 1, 1, 1)$

Given $\varphi: (\mathbb{R}^4, g) \rightarrow \mathbb{R}$, the energy-momentum tensor is defined as

$$T = d\varphi \otimes d\varphi - \frac{1}{2} (g^{\alpha\beta} (d\varphi, d\varphi) - m^2 \varphi^2) g$$

(it's symmetric)

Its components:

$$T_{\alpha\beta} = T_{\beta\alpha}$$

$$T_{\alpha\beta} = \varphi_{,\alpha} \varphi_{,\beta} - \frac{1}{2} g_{\alpha\beta} g^{\gamma\delta} \varphi_{,\gamma} \varphi_{,\delta} - \frac{1}{2} m^2 \varphi^2 g_{\alpha\beta}$$

$$\text{Then } \int_{\text{energy}}^{\alpha} = -T^{\alpha}_{\quad t} \equiv -g^{\alpha\beta} T_{t\beta}$$

$$\int_{\text{energy density}} = T^{tt}$$

$$\int_{\text{momentum}}^{-\alpha} = -T^{\alpha}_{\quad i} \equiv -g^{\alpha\beta} T_{i\beta}, \quad i=1,2,3$$

Momentum density in direction i is $-T^t_i$

Conservation of energy & momentum is encoded in an identity:

$$g^{\beta\gamma} \partial_\gamma T_{\alpha\beta} = \psi_{,\alpha} (\square - m^2) \psi$$

where $\square \psi = g^{\alpha\beta} \psi_{,\alpha\beta}$.

So, if $(\square - m^2) \psi = 0$, then

$$g^{\beta\gamma} \partial_\gamma T_{\alpha\beta} = 0$$

i.e. the divergence of the energy-momentum tensor vanishes

Note: Change of reference frame mixes up energy & momentum. So one often says, "Conservation of energy-momentum", not just one of these.

A symmetry $f: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ is a diffeo s.t.
if $\hat{\varphi} = \varphi \circ f$, then the Lagrangian is preserved.

So if

$$\mathcal{L}(x, \hat{\varphi}, \partial \hat{\varphi}) = \mathcal{L}(x, \varphi, \partial \varphi)$$

then the Lagrangian is preserved. There
is a more general notion of symmetry
called divergence symmetry.

Since, if $\hat{\mathcal{L}} = \mathcal{L} + (\text{div term})$, then

$\mathcal{E}(\hat{\mathcal{L}}) = \mathcal{E}(\mathcal{L})$, diffeos which change the
Lagrangian by a divergence term are also
considered to be symmetries.


We typically study 1-param families of symmetries,

say $F_\lambda: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ } let $\varphi_\lambda \doteq \varphi \circ F_\lambda$. Then

$$\text{we want } \frac{d}{d\lambda} \left(\kappa(\varphi_\lambda, \frac{\partial \varphi_\lambda}{\partial x^k}) \right) = 0.$$

Infinitesimal Symmetry (a vector field) p.55

Space of fields



— path of fields
defined by
continuous
symmetries

$$\delta\varphi \doteq \left. \frac{d\varphi_\lambda}{d\lambda} \right|_{\lambda=0}.$$

This $\delta\varphi$ is an
infinitesimal symmetry.

View it as a vector field
in the space of fields

Since Lagrangian densities which differ by a divergence term gives the same Euler-Lagrange eqns, then we expand our definition of symmetry to also include $f: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ ($\tilde{\varphi} = \varphi \circ f$) st

$$\mathcal{L}(x, \tilde{\varphi}, \partial_x \tilde{\varphi}) = \mathcal{L}(x, \varphi, \partial_x \varphi) + \partial_\alpha V^\alpha.$$

eg. Consider time translation:

$$\varphi(t, x) \mapsto \varphi(t + \lambda, x)$$

Then $\mathcal{L} = -\frac{1}{2} (g^{\alpha\beta} \overbrace{\varphi_{,\alpha} \varphi_{,\beta}}^{\text{means partial derivatives wrt } \alpha \text{ \& } \beta} + m^2 \varphi^2)$

$$g^{\alpha\beta} = \text{diag}(-1, 1, 1, 1)$$

$$\text{Then } \delta \mathcal{L} = -\mathcal{L} g^{\alpha\beta} \varphi_{,\alpha} \dot{\varphi}_{,\beta} + m^2 \varphi \dot{\varphi}$$

$$= \partial_\epsilon \mathcal{L}$$

$$= \partial_\alpha (\delta_\epsilon^\alpha \mathcal{L}). \quad \leftarrow \text{Divergence term}$$

This means there is no preferred instant of time
in KG theory

Noether's Thm

Let $\mathcal{L} = \mathcal{L}(x, \psi, \partial\psi)$. If \mathcal{L} is an infinitesimal
variational symmetry,

then $\delta\mathcal{L} = 0$.

But also, at any pt in the space of fields

$$\delta\mathcal{L} = \mathcal{E}(\mathcal{L})\delta\psi + D_\alpha V^\alpha$$

where $\mathcal{E}(\mathcal{L}) = \frac{\partial\mathcal{L}}{\partial\psi} - D_\alpha\left(\frac{\partial\mathcal{L}}{\partial\psi_{,\alpha}}\right)$

$$\therefore V^\alpha = \frac{\partial\mathcal{L}}{\partial\psi_{,\alpha}}\delta\psi$$

holds for
any field
variation

$$\text{So if } \delta \mathcal{L} = 0, \text{ then } D_\alpha V^\alpha = -\mathcal{E}(\mathcal{L}) \delta \psi$$

This is exactly the type of identity needed for defining a conserved current V^α : if ψ satisfies the KG eqn, then $\mathcal{E}(\mathcal{L}) = 0$ \therefore so

$$D_\alpha V^\alpha = 0.$$

$$\text{Or if } \delta \mathcal{L} = D_\alpha W^\alpha, \text{ then}$$

$$D_\alpha (V^\alpha - W^\alpha) = -\mathcal{E}(\mathcal{L}) \delta \psi$$

\therefore the conserved current is $V^\alpha - W^\alpha$.

Noether's First Thm:

If $\delta\varphi(x, \varphi, \partial\varphi, \dots)$ is a divergence symmetry of $\mathcal{L}(x, \varphi, \partial\varphi)$; i.e.

$$\delta\mathcal{L} = D_\alpha W^\alpha,$$

then there exists a conserved current

given by

$$j^\alpha = \frac{\partial\mathcal{L}}{\partial\varphi_{,\alpha}} \delta\varphi - W^\alpha.$$

Application: the time translation symmetry gives conserved current defining conservation of energy

Recall the time translation is a divergence symmetry:

$$\delta\varphi = \dot{\varphi} \Rightarrow \delta\mathcal{L} = D_\alpha (\delta_t^\alpha \mathcal{L})$$

Recognizing that $\delta\varphi = \vec{\varphi}$, $\delta\mathcal{L} = \delta_t \mathcal{L}$, $\dot{\varphi}$

$$\frac{\partial \mathcal{L}}{\partial \varphi_{,\alpha}} = -g^{\alpha\beta} \varphi_{,\beta}$$

we have: $j^\alpha = -g^{\alpha\beta} \varphi_{,\beta} \varphi_{,t} - \delta_t^\alpha \mathcal{L} = -T_t^\alpha$

This j^α is a conserved current. We did the

calculation $D_\alpha j^\alpha = 0$ earlier when discussing conservation of energy.

We sometimes say the conserved current for energy is

the Noether current associated to time translation symmetry

This was predictable since \mathcal{L} , in the jet space, has no dependence on t , only on $\varphi, d\varphi$.

Space translations & Conservation of Momentum

Let \hat{n} be a const^{unit} vector field on \mathbb{R}^3 . The space

translation is given by $\varphi(t, \vec{x}) \mapsto \varphi_\lambda(t, \vec{x}) \equiv$

$$\vec{x} = (x, y, z) \quad \varphi(t, \vec{x} + \lambda \hat{n})$$

$$\begin{aligned} \text{then } \delta\varphi &\equiv \left. \frac{d\varphi_\lambda}{d\lambda} \right|_{\lambda=0} = \partial_x \varphi(t, \vec{x}) \cdot \hat{n}_x \\ &\quad + \partial_y \varphi(t, \vec{x}) \cdot \hat{n}_y \\ &\quad + \partial_z \varphi(t, \vec{x}) \cdot \hat{n}_z \\ &= \hat{n} \cdot \nabla \varphi_\lambda(t, \vec{x}) \Big|_{\lambda=0} \\ &= \hat{n} \cdot \nabla \varphi \\ &= \hat{n}_i \cdot \varphi_{,i} \end{aligned}$$

To check it is a symmetry, compute

$$\begin{aligned} \delta \mathcal{L} &= \delta \left(\frac{1}{2} \dot{\varphi}^2 - |\nabla \varphi|^2 - m^2 \varphi^2 \right) \\ &= \dot{\varphi} \delta \dot{\varphi} - \nabla \varphi \cdot \nabla \delta \varphi - m^2 \varphi \cdot \delta \varphi \\ &= \dot{\varphi} (\hat{n} \cdot \nabla \dot{\varphi}) - \nabla \varphi \cdot \nabla (\hat{n} \cdot \nabla \varphi) - m^2 \varphi (\hat{n} \cdot \nabla \varphi) \\ &= \hat{n} \left(\dot{\varphi} \cdot \nabla \dot{\varphi} - \nabla \varphi \cdot \nabla^2 \varphi - m^2 \varphi \cdot \nabla \varphi \right) \end{aligned}$$

\hat{n} is
const

On the other hand:

$$\nabla \mathcal{L} = \dot{\psi} \nabla \dot{\psi} - \nabla \psi \cdot \nabla^2 \psi - m^2 \psi \nabla \psi$$

$$\text{So } \delta \mathcal{L} = \hat{n} \cdot \nabla \mathcal{L} = \nabla (\hat{n} \cdot \mathcal{L})$$

$$= \underbrace{D_\alpha}_{\text{total derivative}} (W^\alpha) \quad \text{where} \\ W^\alpha = (0, n^i \mathcal{L})$$

So space translation = divergence symmetry.

$$\text{Then, } j^\alpha = (p, j^i) = \frac{\partial \mathcal{L}}{\partial \psi, \alpha} \delta \psi - W^\alpha \quad \text{by Noether's theorem}$$

$$\text{So } p = \dot{\psi} \delta \psi - 0 = \dot{\psi} \hat{n} \cdot \nabla \psi.$$

$$\text{Since } |\nabla \psi|^2 = \psi_{,1}^2 + \psi_{,2}^2 + \psi_{,3}^2, \text{ then}$$

$$j^i = -\psi_{,i} \hat{n} \cdot \nabla \psi - \frac{n^i}{2} \mathcal{L}$$

$$p = \dot{\psi} \hat{n} \cdot \nabla \psi, \quad j^i = -\psi_i \hat{n} \cdot \nabla \psi + \frac{1}{2} \hat{n}^i (|\nabla \psi|^2 - \dot{\psi}^2 + m^2 \psi^2)$$

Let's verify $\frac{dp}{dt} + \nabla \cdot \vec{j} = 0$.

$$\frac{dp}{dt} = \ddot{\psi} \hat{n} \cdot \nabla \psi + \dot{\psi} \hat{n} \cdot \nabla \dot{\psi}$$

once summed over i , these cancel.

$$\partial_i j^i = -\psi_{;i} \hat{n} \cdot \nabla \psi - \psi_i \hat{n} \cdot \nabla \psi_i + \hat{n}^i (\nabla \psi_i \cdot \nabla \psi - \dot{\psi} \cdot \dot{\psi}_i + m^2 \psi \cdot \psi_i)$$

$$\Rightarrow \nabla \cdot \vec{j} = -\nabla^2 \psi \cdot \hat{n} \cdot \nabla \psi - \dot{\psi} \hat{n} \cdot \nabla \dot{\psi} + m^2 \psi \cdot \hat{n} \cdot \nabla \psi$$

$$\frac{dp}{dt} + \nabla \cdot \vec{j} = \ddot{\psi} \hat{n} \cdot \nabla \psi - \nabla^2 \psi \cdot \hat{n} \cdot \nabla \psi + m^2 \psi \hat{n} \cdot \nabla \psi$$

$$= -(\square - m^2)\psi \cdot \hat{n} \cdot \nabla \psi = 0$$

" 0 since ψ is a KG field.

Since \hat{n} is arbitrary, we actually have 3 indep conservation laws for 3 (nearly) indep choices of \hat{n} .

Angular momentum.

Q: What symmetries conserve angular momentum?

A: Lorentz symmetries.

def. A Lorentz symmetry of \mathbb{R}^4 is a linear transform

$$x^\alpha \mapsto S^\alpha_\beta x^\beta$$

s.t. the quadratic form $g_{\alpha\beta} x^\alpha x^\beta = -t^2 + x^2 + y^2 + z^2$
is invariant.

$$S^\alpha_\gamma S^\beta_\delta g_{\alpha\beta} = g_{\gamma\delta} \quad (*)$$

let $S(\lambda)$ be a 1-param family of Lorentz symm.

$$\text{s.t. } S^\alpha_\beta(0) = \delta^\alpha_\beta, \quad \omega^\alpha_\beta \equiv \left. \frac{\partial S^\alpha_\beta}{\partial \lambda} \right|_{\lambda=0}$$

Differentiating $(*)$ w.r.t λ we get

$$\omega^\alpha_\gamma g_{\alpha\delta} + \omega^\beta_\delta g_{\alpha\beta} = 0$$

Define $\omega_{\alpha\beta} = g_{\beta\gamma} \omega_{\alpha}^{\gamma}$. Then Lorentz transf
are generated by ω iff $\omega_{\alpha\beta} = -\omega_{\beta\alpha}$.

So the Lie algebra consists of 4×4 matrices X
s.t. $g X g = -X^T$ ($g = \text{diag}(-1, 1, 1, 1)$).

$$\text{Then } \delta\varphi = (\omega_{\beta}^{\alpha} x^{\beta}) \varphi_{,\alpha}$$

$$\delta\mathbb{L} = D_{\alpha} (\omega_{\beta}^{\alpha} x^{\beta} \mathbb{L})$$

$$\} \quad \tilde{j}^{\alpha} = \omega_{\beta\gamma} M^{\alpha(\beta)(\gamma)}$$

conserved currents associated to
relativistic angular momentum

Class:

All continuous isometries of flat spacetime are contained in
the Poincaré group = diffeomorphisms generated by Lorentz
transformations & spacetime translations.

Internal Symmetries

These are symmetries on the space of fields, not on spacetime. Eg. $\psi \rightarrow -\psi$.

Here, unless $m=0$, there are no interesting continuous internal symmetries.

eg. $m=0$, $\psi \rightarrow \psi + \lambda$

$$\text{Then } \delta\psi = 1 \quad \delta\mathcal{L} = \dot{\psi} \delta\dot{\psi} - \nabla\psi \cdot \nabla\delta\psi$$

$$\delta\dot{\psi} = \left. \frac{d\dot{\psi}_\lambda}{d\lambda} \right|_{\lambda=0} = 0 \quad \text{" } D_\alpha \psi^\alpha$$

$$\text{So } \vec{j}^\alpha = \frac{\partial\mathcal{L}}{\partial\psi_{,\alpha}} \delta\psi - \underbrace{W^\alpha}_{\text{const?}}$$

$$\text{Seems } \rho = \dot{\psi} \quad \Rightarrow \quad \frac{d\rho}{dt} + \nabla \cdot \vec{j} = -\square\psi = 0$$

$\vec{j}^\alpha = -\psi_{,\alpha}$

Since $m=0$ }

$$\text{So } \square\psi = 0.$$

Charged KG Field & its internal symmetries.

A charged KG Field is a \mathbb{C} -valued ϕ^n

$$\phi: M \rightarrow \mathbb{C} \quad g_{\alpha\beta} = g^{\alpha\beta} = \text{diag}(-1, 1, 1, 1)$$

The Lagrangian is now $\mathcal{L} = -g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi^* + m^2 |\phi|^2$

We can write this Lagrangian as the sum of two \mathbb{R} -valued Lagrangians w/ masses $m_1 = m_2 = m$
} fields ϕ_1, ϕ_2 .

Then the charged field can be written as

$$\phi = \frac{1}{\sqrt{2}} (\phi_1 + i \phi_2)$$

So really, we can take in real & imaginary parts if we want: ϕ & ϕ^*

The Euler-Lagrange eqns are now:

$$\mathcal{E}_\psi(\mathcal{L}) = (\square - m^2)\psi^* = 0$$

$$\mathcal{E}_{\psi^*}(\mathcal{L}) = (\square - m^2)\psi = 0$$

Working on \mathbb{C} gives new deg of freedom which allows us to introduce conserved electric charge.

In QFT, we get anti-particles.

The charged Lagrangian now admits the internal continuous symmetry $\psi_t = e^{it} \psi$, $\psi_t^* = e^{-it} \psi^*$.

These are sometimes called phase transformations or rigid $U(1)$ transformations.

See p. 66-7 for the conserved current - the answer is

$$j^\alpha = -ig^{\alpha\beta} (\psi^* \psi_{,\beta} - \psi_{,\beta} \psi^*)$$

Let $V \subset \mathbb{R}^3$ at a fixed time. Then the total U(1) charge

$$Q = i \int_V (\psi^* \dot{\psi} - \dot{\psi} \psi^*) d^3x$$

Use this for modeling electric charge in electrodynamics

Can also use it to model charge-current interactions w/ neutral currents in electroweak theory.

Of course, we can do all this for general groups w/ representations on vect space V .

Consider $\psi: M \rightarrow V$ w/ internal symmetries given by $\rho: G \rightarrow \text{Aut}(V)$.

Or even, let V be a vector bundle & $\psi: M \rightarrow V$ a section

$$\text{Let } G = SU(2), \quad V = \mathbb{C}^2.$$

an element $U \in SU(2)$ can be written as

$$U(\theta, n) = \cos\left(\frac{\theta}{2}\right) I + i \sin\left(\frac{\theta}{2}\right) n^i \sigma_i$$

$$n = (n^1, n^2, n^3), \quad \sum_{j=1}^3 (n^j)^2 = 1 \quad \zeta$$

Pauli
↓
matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

IF $U(\lambda)$ is any 1-param family in $SU(2)$, w/ $U(0) = I$

$$\zeta \quad \psi_\lambda = U(\lambda)\psi, \text{ then}$$

$$S\psi = i\tau\psi, \text{ where } \tau \text{ is a Hermitian, traceless}$$

2×2
matrix

$$\text{defined by } \tau = \frac{1}{i} \begin{pmatrix} \frac{dU}{d\lambda} \\ \frac{dU}{d\lambda} \end{pmatrix}_{\lambda=0}$$

$$\text{So } i\tau \in SU(2).$$

Of course, τ is a linear combo of the σ_i

Hermitian inner prod on \mathbb{C}^2 : $(\varphi_1, \varphi_2) = \varphi_1^* \varphi_2$

Of course, it is $U(2)$ invariant. So we build a Lag.

$$\mathcal{L} = - \left[g^{\alpha\beta} (\varphi_{,\alpha}, \varphi_{,\beta}) + m^2 (\varphi, \varphi) \right]$$

Claim: For symmetry $\delta\varphi = i\epsilon\delta$, the assoc conserved

$$\text{current is } j^\alpha = i g^{\alpha\beta} (\varphi_{,\beta}^* \tau \varphi - \varphi^* \tau \varphi_{,\beta})$$

There are 3 indep conserved currents here b/c of τ .

The 3 conserved charges assoc w/ $SU(2)$ symm are

called isospin

Claim: The converse to what we wrote as

Noether's 1st Thm is true:

For each conservation law for a system of

Euler-Lagrange eqns, there is a corresponding
symmetry of the Lagrangian.

For many types of field theories, we also have

1:1 correspondence b/w conservation laws &
symmetries of the Lagrangian.

(including KG field theory)

Let $M = (\mathbb{R}^{2n}, \omega)$ (phase space) w/ $H: \mathbb{R}^{2n} \rightarrow \mathbb{R}$
a Hamiltonian. Let $\gamma: [0,1] \rightarrow \mathbb{R}^{2n}$ be a classical traj.

Then for any function F , $\{F, H\} = \frac{d}{dt} F(\gamma(t))$.

$$\begin{aligned} \text{This is b/c we define } \{F, H\} &\equiv -(\nabla F)^T \mathcal{J}_0 \nabla H \\ &= (\nabla F)^T \dot{\gamma} \\ &= \frac{d}{dt} (F \circ \gamma) \end{aligned}$$

More generally, since $\{, \dot{\gamma}\}: C^\infty(M) \times C^\infty(M) \rightarrow C^\infty(M)$,

take another function G ; we can think of
 G as the infinitesimal generator of a transformation:

$$\delta F = \{F, G\}$$

Then, if $\delta H = \{H, G\} = 0$, we call G a symmetry
of the Hamiltonian system.

$$\text{But } \dot{G} = \{G, H\} = -\{H, G\} = 0$$

So G is also a conserved quantity along class. traj.

i.e. A F^n on phase space generates a

Symmetry iff it is conserved.